

113-2 Semester General Chemistry Midterm Exam(C)-20250409

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

1) Which of the following could be added to a solution of acetic acid to prepare a buffer?

- A) sodium hydroxide only
- B) sodium acetate only
- C) hydrofluoric acid or nitric acid
- D) sodium acetate or sodium hydroxide
- E) nitric acid only

Answer: D

2) Calculate the pH of a solution that is 0.310 M in sodium formate (NaHCO_2) and 0.190 M in formic acid (HCO_2H). The K_a of formic acid is 1.77×10^{-4} .

- A) 13.79
- B) 3.532
- C) 10.04
- D) 4.975
- E) 3.958


Answer: E

3) Calculate the percent ionization of formic acid (HCO_2H) in a solution that is 0.152 M in formic acid. The K_a of formic acid is 1.77×10^{-4} .

- A) 3.44
- B) 0.0180
- C) 2.74×10^{-5}
- D) 0.581
- E) 8.44

Answer: A

4)



A 25.0 mL sample of a solution of an unknown compound is titrated with a 0.115 M NaOH solution. The titration curve above was obtained. The unknown compound is _____.

- A) a weak acid
- B) a strong base
- C) a weak base
- D) a strong acid
- E) neither an acid nor a base

Answer: A

5) A solution of NaF is added dropwise to a solution that is 0.0144 M in Ba^{2+} . When the concentration of F^- exceeds _____ M, BaF_2 will precipitate. Neglect volume changes. For BaF_2 , $K_{sp} = 1.7 \times 10^{-6}$.

- A) 1.1×10^{-2}
- B) 2.4×10^{-8}
- C) 5.9×10^{-5}
- D) 1.2×10^{-4}
- E) 2.7×10^{-3}

Answer: A

6) What is the molar solubility of manganese carbonate (MnCO_3) in water? The solubility-product constant for MnCO_3 is 5.0×10^{-10} at 25 °C.

- A) 3.2×10^{-5}
- B) 9.30
- C) 2.2×10^{-5}
- D) 1.0×10^{-9}
- E) 2.5×10^{-10}

Answer: C

7) Consider the following table of K_{sp} values.

Name	Formula	K_{sp}
Cadmium carbonate	CdCO_3	5.2×10^{-12}
Cadmium hydroxide	Cd(OH)_2	2.5×10^{-14}
Calcium fluoride	CaF_2	3.9×10^{-11}
Silver iodide	AgI	8.3×10^{-17}
Zinc carbonate	ZnCO_3	1.4×10^{-11}

Which compound listed below has the greatest molar solubility in water?

- A) CdCO_3 B) CaF_2 C) ZnCO_3 D) Cd(OH)_2 E) AgI

Answer: B

8) A 25.0 mL sample of 0.723 M HClO_4 is titrated with a 0.273 M KOH solution. The H_3O^+ concentration after the addition of 50.0 mL of KOH is _____ M.

- A) 0.0587
B) 0.273
C) 0.430
D) 0.0181
E) none of the above

Answer: A

9) Which one of the following is least soluble in water?

- A) $\text{CH}_3\text{CH}_2\text{OH}$
B) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$
C) CH_3OH
D) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
E) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$

Answer: B

10) Calculate the molality of a 10.0% (by mass) aqueous solution of hydrochloric acid (HCl).

- A) 2.74 m
B) 3.05 m
C) 0.274 m
D) 4.33 m
E) The density of the solution is needed to solve the problem.

Answer: B

11) Calculate the mole fraction of nitric acid of a 17.5% (by mass) aqueous solution of nitric acid (HNO_3).

- A) 0.278 B) 3.37 C) 0.0607 D) 0.0572 E) 1.75

Answer: D

- 12) The concentration of CO_2 in a soft drink bottled with a partial pressure of CO_2 of 4.0 atm over the liquid at 25 °C is 1.2×10^{-1} M. The Henry's law constant for CO_2 at this temperature is _____.
- A) 3.0×10^{-2} mol/L-atm
 - B) 4.5×10^{-3} mol/L-atm
 - C) 2.3×10^{-2} mol/L-atm
 - D) 5.6×10^{-3} mol/L-atm
 - E) More information is needed to solve the problem.

Answer: A

- 13) A 1.35 m aqueous solution of compound X had a boiling point of 101.4 °C. Which one of the following could be compound X? The boiling point elevation constant for water is 0.52 °C/m.
- A) Na_3PO_4
 - B) CaCl_2
 - C) $\text{C}_6\text{H}_{12}\text{O}_6$
 - D) KCl
 - E) $\text{CH}_3\text{CH}_2\text{OH}$

Answer: D

- 14) A solution contains 15 ppm of benzene. The density of the solution is 1.00 g/mL. This means that _____.
- A) the solution is 15% by mass of benzene
 - B) there are 15 mg of benzene in 1.0 g of this solution
 - C) 1.0 g of the solution contains 15×10^{-6} g of benzene
 - D) 100 g of the solution contains 15 g of benzene
 - E) 1.0 L of the solution contains 15 g of benzene

Answer: C


- 15) The osmotic pressure of a solution formed by dissolving 80.0 mg of aspirin ($\text{C}_9\text{H}_8\text{O}_4$) in 0.250 L of water at 25 °C is _____ atm. ($R = 0.08206$ L-atm/K-mol)
- A) 43.5
 - B) 4.41
 - C) 0.0435
 - D) 7.83
 - E) 3.65×10^{-3}

Answer: C

- 16) A solution is prepared by dissolving 24.7 g of CaCl_2 in 375 g of water. The density of the resulting solution is 1.05 g/mL. The concentration of CaCl_2 is _____% by mass.
- A) 6.49
 - B) 0.0618
 - C) 0.0649
 - D) 6.18
 - E) 6.24

Answer: D

- 17) At elevated temperatures, dinitrogen pentoxide decomposes to nitrogen dioxide and oxygen:




When the rate of formation of O_2 is 2.2×10^{-4} M/s, the rate of decomposition of N_2O_5 is _____ M/s.

- A) 2.8×10^{-4}
- B) 1.1×10^{-4}
- C) 5.5×10^{-4}
- D) 4.4×10^{-4}
- E) 2.2×10^{-4}

Answer: D

18) - 19) The data in the table below were obtained for the reaction:



Experiment Number	$[\text{ClO}_2] (\text{M})$	$[\text{OH}^-] (\text{M})$	Initial Rate (M/s)
1	0.060	0.030	0.0248
2	0.020	0.030	0.00276
3	0.020	0.090	0.00828

18) What is the order of the reaction with respect to ClO_2 ?

- A) 0 B) 3 C) 2 D) 1 E) 4


Answer: C

19) What is the magnitude of the rate constant for the reaction?

- A) 713 B) 230 C) 115 D) 1.15×10^4 E) 4.6

Answer: B

20) The reaction



At 230.3°C , $k = 6.29 \times 10^{-4} \text{ s}^{-1}$. If $[\text{CH}_3\text{-N}\equiv\text{C}]$ is 1.00×10^{-3} initially, $[\text{CH}_3\text{-N}\equiv\text{C}]$ is _____ after $1.000 \times 10^3 \text{ s}$.

- A) 2.34×10^{-4} B) 4.27×10^{-3} C) 1.88×10^{-3} D) 5.33×10^{-4} E) 1.00×10^{-6}


Answer: D

21) A compound decomposes by a first-order process. If 25.0% of the compound decomposes in 60.0 minutes, the half-life of the compound is _____.

- A) 65 minutes B) 120 minutes C) 145 minutes D) 198 minutes E) 180 minutes

Answer: C


22) Which energy difference in the energy profile below corresponds to the activation energy for the forward reaction?




- A) x B) y C) x + y D) y - x E) x - y

Answer: A

23) A possible mechanism for the overall reaction



is



The rate law for formation of NOBr based on this mechanism is rate = _____.

- A) $k_1[\text{NO}]^{1/2}$
- B) $(k_2k_1/k_{-1})[\text{NO}][\text{Br}_2]^2$
- C) $k_1[\text{Br}_2]^{1/2}$
- D) $(k_1/k_{-1})^2[\text{NO}]^2$
- E) $(k_2k_1/k_{-1})[\text{NO}]^2[\text{Br}_2]$

Answer: E

24) A particular first-order reaction has a rate constant of $1.35 \times 10^2 \text{ s}^{-1}$ at 25.0°C . What is the magnitude of k at 75.0°C if $E_a = 60.2 \text{ kJ/mol}$? ($R = 8.3145 \text{ J/K}\cdot\text{mol}$)

- A) 4.43×10^3
- B) 1.35×10^2
- C) 2.71×10^6
- D) 2.44×10^4
- E) 471


Answer: A

25) At equilibrium, _____.


- A) the rate constants of the forward and reverse reactions are equal
- B) all chemical reactions have ceased
- C) the rates of the forward and reverse reactions are equal
- D) the value of the equilibrium constant is 1
- E) the limiting reagent has been consumed

Answer: C

26) The value of K_{eq} for the following reaction is 0.25:




The value of K_{eq} at the same temperature for the reaction below is _____.



- A) 16
- B) 0.50
- C) 0.063
- D) 0.12
- E) 0.25

Answer: C


27) Which of the following expressions is the correct equilibrium-constant expression for the reaction below?



- A) $1 / [(\text{NH}_4)_2\text{Se}]$
- B) $[\text{NH}_3][\text{H}_2\text{Se}] / [(\text{NH}_4)_2\text{Se}]$
- C) $[\text{NH}_3]^2[\text{H}_2\text{Se}]$
- D) $[(\text{NH}_4)_2\text{Se}] / [\text{NH}_3]^2[\text{H}_2\text{Se}]$
- E) $[\text{NH}_3]^2[\text{H}_2\text{Se}] / [(\text{NH}_4)_2\text{Se}]$

Answer: C

28) Consider the following reaction at equilibrium:




Le Châtelier's principle predicts that adding $\text{O}_2 \text{ (g)}$ to the reaction container will _____.

- A) increase the partial pressure of $\text{CO}_2 \text{ (g)}$ at equilibrium
- B) increase the value of the equilibrium constant
- C) decrease the partial pressure of $\text{CO}_2 \text{ (g)}$ at equilibrium
- D) decrease the value of the equilibrium constant
- E) increase the partial pressure of CO (g) at equilibrium

Answer: A

29) Phosphorous trichloride and phosphorous pentachloride equilibrate in the presence of molecular chlorine according to the reaction:



An equilibrium mixture at 450 K contains

$\text{PCl}_3 = 0.224 \text{ atm}$,


$\text{PCl}_2 = 0.284 \text{ atm}$, and

$\text{PCl}_5 = 4.24 \text{ atm}$. What is the value of K_p at this temperature?

- A) 1.50×10^{-2}
- B) 2.70×10^{-1}
- C) 3.74
- D) 66.7
- E) 8.36

Answer: D

30) Dinitrogen tetroxide partially decomposes according to the following equilibrium:




A 1.000-L flask is charged with $9.20 \times 10^{-3} \text{ mol}$ of N_2O_4 . At equilibrium, $5.98 \times 10^{-3} \text{ mol}$ of N_2O_4 remains. K_{eq} for this reaction is _____.

- A) 0.183
- B) 2.96×10^{-5}
- C) 6.94×10^{-3}
- D) 0.197
- E) 0.212

Answer: C


31) Given the following reaction at equilibrium, if $K_c = 5.84 \times 10^5$ at 230.0°C , $K_p =$ _____.



- A) 6.44×10^5 B) 3.67×10^{-2} C) 1.41×10^4 D) 2.41×10^7 E) 2.40×10^6

Answer: C

32) Consider the following reaction at equilibrium:



Le Châtelier's principle predicts that the moles of H_2 in the reaction container will increase with _____.

- A) a decrease in the total volume of the reaction vessel (T constant)
B) a decrease in the total pressure (T constant)
C) some removal of NH_3 from the reaction vessel (V and T constant)
D) an increase in total pressure by the addition of helium gas (V and T constant)
E) addition of some N_2 to the reaction vessel (V and T constant)

Answer: B

33) A Brønsted-Lowry base is defined as a substance that _____.

- A) increases $[\text{H}^+]$ when placed in H_2O
B) acts as a proton donor
C) increases $[\text{OH}^-]$ when placed in H_2O
D) acts as a proton acceptor
E) decreases $[\text{H}^+]$ when placed in H_2O

Answer: D

34) Of the acids in the table below, _____ is the strongest acid.

Acid	K_a
HOAc	1.8×10^{-5}
HCHO_2	1.8×10^{-4}
HClO	3.0×10^{-8}
HF	6.8×10^{-4}

- A) HOAc
B) HF
C) HCHO_2
D) HClO
E) HOAc and HCHO_2

Answer: B

35) The pH of an aqueous solution at 25.0°C is 10.40. What is the molarity of H^+ in this solution?

- A) 2.5×10^{-4} B) 1.0×10^{-13} C) 3.60 D) 2.5×10^{10} E) 4.0×10^{-11}

Answer: E

36) A $8.0 \times 10^{-3} \text{ M}$ aqueous solution of Ca(OH)_2 at 25.0°C has a pH of _____.

- A) 6.3×10^{-13} B) 1.6×10^{-2} C) 11.90 D) 1.80 E) 12.20

Answer: E

37) The K_a of hypochlorous acid (HClO) is 3.0×10^{-8} at 25.0°C . Calculate the pH of a 0.0335 M hypochlorous acid solution.

A) 3.02

B) 4.50

C) -3.02

D) 9.50

E) 6.52

Answer: B

38) Calculate the pH of a 0.250 M aqueous solution of NH_3 . The K_b of NH_3 is 1.77×10^{-5} .

A) 2.08

B) 11.32

C) 2.68

D) 11.92

E) 8.95

Answer: B

39) K_b for NH_3 is 1.8×10^{-5} . What is the pH of a 0.35 M aqueous solution of NH_4Cl at 25.0°C ?

A) 11.23

B) 4.85

C) 2.60

D) 9.15

E) 11.40

Answer: B

40) Of the following, which is the strongest acid?

A) HClO_4

B) HClO_2

C) HIO

D) HClO_3

E) HClO

Answer: A