


114-1 Semester General Chemistry Midterm Exam ^A~~(B)~~-20251105

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

- 1) When the following equation is balanced, the coefficients are _____.



- A) 2, 12, 8, 9 B) 4, 4, 32, 36 C) 1, 4, 8, 9 D) 2, 3, 4, 4 E) 2, 25, 16, 18

Answer: E

- 2) Which one of the following is not true concerning automotive air bags?

- A) They are loaded with sodium azide initially
 B) A gas is produced when the air bag activates.
 C) They are inflated as a result of a decomposition reaction
 D) The gas used for inflating them is oxygen
 E) The two products of the decomposition reaction are sodium and nitrogen

Answer: D

- 3) Which of the following are decomposition reactions?

- 1) $\text{CH}_4(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
 2) $\text{CaO}(\text{s}) + \text{CO}_2(\text{g}) \rightarrow \text{CaCO}_3(\text{s})$
 3) $\text{Mg}(\text{s}) + \text{O}_2(\text{g}) \rightarrow \text{MgO}(\text{s})$
 4) $\text{PbCO}_3(\text{s}) \rightarrow \text{PbO}(\text{s}) + \text{CO}_2(\text{g})$

- A) 4 only B) 1, 2, 3, and 4 C) 2, 3, and 4 D) 2 and 3 E) 1, 2, and 3

Answer: A

- 4) What is the mass % of carbon in dimethylsulfoxide ($\text{C}_2\text{H}_6\text{SO}$) rounded to three significant figures?

- A) 60.0 B) 7.74 C) 30.7 D) 20.6 E) 79.8

Answer: C

- 5) Gaseous argon has a density of 1.40 g/L at standard conditions. How many argon atoms are in 1.00 L of argon gas at standard conditions?

- A) 3.43×10^{25} B) 6.02×10^{23} C) 1.59×10^{25} D) 2.11×10^{22} E) 4.76×10^{22}

Answer: D

- 6) The total number of atoms in 0.111 mol of $\text{Fe}(\text{CO})_3(\text{PH}_3)_2$ is _____.

- A) 15.0 B) 4.46×10^{21} C) 2.76×10^{-24} D) 1.00×10^{24} E) 1.67

Answer: D


- 7) The molecular formula of aspartame, the generic name of NutraSweet[®], is $\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}_5$.

The molar mass of aspartame, rounded to the nearest integer, is _____ g.

- A) 43 B) 24 C) 39 D) 156 E) 294

Answer: E

- 8) Sulfur and oxygen react to produce sulfur trioxide. In a particular experiment, 7.9 grams of SO_3 are produced by the reaction of 5.0 grams of O_2 with 6.0 grams of S. What is the % yield of SO_3 in this experiment?



- A) 95 B) 99 C) 63 D) 32 E) 75

Answer: A

- 9) The net ionic equation for formation of an aqueous solution of NiI_2 accompanied by evolution of CO_2 gas via mixing solid NiCO_3 and aqueous hydriodic acid is _____.

- A) $2\text{NiCO}_3 \text{ (s)} + \text{HI (aq)} \rightarrow 2\text{H}_2\text{O (l)} + \text{CO}_2 \text{ (g)} + 2\text{Ni}^{2+} \text{ (aq)}$
B) $\text{NiCO}_3 \text{ (s)} + 2\text{HI (aq)} \rightarrow \text{H}_2\text{O (l)} + \text{CO}_2 \text{ (g)} + \text{Ni}^{2+} \text{ (aq)} + 2\text{I}^- \text{ (aq)}$
C) $\text{NiCO}_3 \text{ (s)} + 2\text{HI (aq)} \rightarrow 2\text{H}_2\text{O (l)} + \text{CO}_2 \text{ (g)} + \text{NiI}_2 \text{ (aq)}$
D) $\text{NiCO}_3 \text{ (s)} + \text{I}^- \text{ (aq)} \rightarrow 2\text{H}_2\text{O (l)} + \text{CO}_2 \text{ (g)} + \text{Ni}^{2+} \text{ (aq)} + \text{HI (aq)}$
E) $\text{NiCO}_3 \text{ (s)} + 2\text{H}^+ \text{ (aq)} \rightarrow \text{H}_2\text{O (l)} + \text{CO}_2 \text{ (g)} + \text{Ni}^{2+} \text{ (aq)}$

Answer: E

- 10) When aqueous solutions of _____ are mixed, a precipitate forms.

- A) NaI and KBr
B) Li_2CO_3 and CsI
C) KOH and $\text{Ba}(\text{NO}_3)_2$
D) K_2SO_4 and CrCl_3
E) NiBr_2 and AgNO_3

Answer: E

- 11) Which combination will produce a precipitate?

- A) $\text{NH}_4\text{OH (aq)}$ and HCl (aq)
B) NaOH (aq) and $\text{Fe}(\text{NO}_3)_2 \text{ (aq)}$
C) NaCl (aq) and $\text{HC}_2\text{H}_3\text{O}_2 \text{ (aq)}$
D) $\text{AgNO}_3 \text{ (aq)}$ and $\text{Ca}(\text{C}_2\text{H}_3\text{O}_2)_2 \text{ (aq)}$
E) NaOH (aq) and HCl (aq)

Answer: B

- 12) Which one of the following is not true concerning 2.00 L of 0.100 M solution of $\text{Ca}_3(\text{PO}_4)_2$?

- A) This solution contains 1.60 mol of oxygen atoms.
B) 1.00 L of this solution is required to furnish 0.300 mol of Ca^{2+} ions.
C) There are 6.02×10^{22} phosphorus atoms in 500.0 mL of this solution.
D) This solution contains 0.200 mol of $\text{Ca}_3(\text{PO}_4)_2$.
E) This solution contains 6.67×10^{-2} mol of Ca^{2+} .

Answer: A

Answer : E

- 13) What mass (g) of CaF_2 is formed when 47.8 mL of 0.334 M NaF is treated with an excess of aqueous calcium nitrate?

- A) 0.623 B) 2.49 C) 0.472 D) 0.943 E) 1.25

Answer: A

14) What volume (mL) of 7.48×10^{-2} M perchloric acid can be neutralized with 115 mL of 0.244 M sodium hydroxide?

- A) 8.60 B) 125 C) 188 D) 375 E) 750

Answer: D

15) The molarity (M) of an aqueous solution containing 22.5 g of sucrose ($C_{12}H_{22}O_{11}$) in 35.5 mL of solution is _____.

- A) 1.85 B) 0.104 C) 1.85×10^{-3} D) 0.0657 E) 3.52

Answer: A

16) Based on the activity series, which one of the reactions below will occur

TABLE 4.5 Activity Series of Metals in Aqueous Solution

Metal	Oxidation Reaction
Lithium	$Li(s) \longrightarrow Li^+(aq) + e^-$
Potassium	$K(s) \longrightarrow K^+(aq) + e^-$
Barium	$Ba(s) \longrightarrow Ba^{2+}(aq) + 2e^-$
Calcium	$Ca(s) \longrightarrow Ca^{2+}(aq) + 2e^-$
Sodium	$Na(s) \longrightarrow Na^+(aq) + e^-$
Magnesium	$Mg(s) \longrightarrow Mg^{2+}(aq) + 2e^-$
Aluminum	$Al(s) \longrightarrow Al^{3+}(aq) + 3e^-$
Manganese	$Mn(s) \longrightarrow Mn^{2+}(aq) + 2e^-$
Zinc	$Zn(s) \longrightarrow Zn^{2+}(aq) + 2e^-$
Chromium	$Cr(s) \longrightarrow Cr^{3+}(aq) + 3e^-$
Iron	$Fe(s) \longrightarrow Fe^{2+}(aq) + 2e^-$
Cobalt	$Co(s) \longrightarrow Co^{2+}(aq) + 2e^-$
Nickel	$Ni(s) \longrightarrow Ni^{2+}(aq) + 2e^-$
Tin	$Sn(s) \longrightarrow Sn^{2+}(aq) + 2e^-$
Lead	$Pb(s) \longrightarrow Pb^{2+}(aq) + 2e^-$
Hydrogen	$H_2(g) \longrightarrow 2H^+(aq) + 2e^-$
Copper	$Cu(s) \longrightarrow Cu^{2+}(aq) + 2e^-$
Silver	$Ag(s) \longrightarrow Ag^+(aq) + e^-$
Mercury	$Hg(l) \longrightarrow Hg^{2+}(aq) + 2e^-$
Platinum	$Pt(s) \longrightarrow Pt^{2+}(aq) + 2e^-$
Gold	$Au(s) \longrightarrow Au^{3+}(aq) + 3e^-$



- A) $Fe(s) + ZnCl_2(aq) \rightarrow FeCl_2(aq) + Zn(s)$
 B) $Pb(s) + NiI_2(aq) \rightarrow PbI_2(aq) + Ni(s)$
 C) $SnBr_2(aq) + Cu(s) \rightarrow CuBr_2(aq) + Sn(s)$
 D) $Mn(s) + NiCl_2(aq) \rightarrow MnCl_2(aq) + Ni(s)$
 E) None of the reactions will occur.

Answer: D

17) Which one of the following is an exothermic process?

- A) condensation of water vapor
- B) ice melting
- C) boiling soup
- D) water evaporating
- E) Ammonium thiocyanate and barium hydroxide are mixed at 25 °C: the temperature drops.

Answer: A

18) Which one of the following statements is true?

- A) Enthalpy is a state function.
- B) H is the value of q measured under conditions of constant volume.
- C) The enthalpy change of a reaction is the reciprocal of the ΔH of the reverse reaction.
- D) The enthalpy change for a reaction is independent of the state of the reactants and products.
- E) Enthalpy is an intensive property.

Answer: A

19) The temperature of a 12.58 g sample of calcium carbonate [CaCO_3 (s)] increases from 23.6 °C to 38.2 °C. If the specific heat of calcium carbonate is 0.82 J/g·K, how many joules of heat are absorbed?

- A) 5.0
- B) 151
- C) 410
- D) 7.5
- E) 0.82

Answer: B

20) For which one of the following equations is $\Delta H^\circ_{\text{rxn}}$ equal to ΔH°_f for the product?

- A) $\text{Xe (g)} + 2\text{F}_2 \text{ (g)} \rightarrow \text{XeF}_4 \text{ (g)}$
- B) $\text{CH}_4 \text{ (g)} + 2\text{Cl}_2 \text{ (g)} \rightarrow \text{CH}_2\text{Cl}_2 \text{ (l)} + 2\text{HCl (g)}$
- C) $2\text{CO (g)} + \text{O}_2 \text{ (g)} \rightarrow 2\text{CO}_2 \text{ (g)}$
- D) $\text{C (diamond)} + \text{O}_2 \text{ (g)} \rightarrow \text{CO}_2 \text{ (g)}$
- E) $\text{N}_2 \text{ (g)} + \text{O}_3 \text{ (g)} \rightarrow \text{N}_2\text{O}_3 \text{ (g)}$


Answer: A

21) The change in the internal energy of a system that absorbs 2,500 J of heat and that does 7,655 J of work on the surroundings is _____ J.

- A) 5,155
- B) -5,155
- C) -10,155
- D) 1.91×10^7
- E) 10,155

Answer: B

22) The value of ΔH° for the reaction below is -790 kJ. The enthalpy change accompanying the reaction of 0.95 g of S is _____ kJ.



- A) 23
- B) -790
- C) 12
- D) -23
- E) -12

Answer: E

23) The value of ΔH° for the reaction below is +128.1 kJ:




How many kJ of heat are consumed when 5.10 g of H_2 (g) is formed as shown in the equation?



- A) 326 B) 62.0 C) 653 D) 163 E) 128

Answer: D

24) ΔH for the reaction



is _____ kJ, give the data below.



- A) +1135 B) +355 C) -35 D) -1135 E) +35

Answer: B

25) The specific heat of liquid bromine is 0.226 J/g-K. How much heat (J) is required to raise the temperature of 10.0 mL of bromine from 25.00 °C to 27.30 °C? The density of liquid bromine: 3.12 g/mL.

- A) 32.4 B) 5.20 C) 10.4 D) 16.2 E) 300

Answer: D

26) All of the orbitals in a given subshell have the same value of the _____ quantum number.

- A) angular momentum
B) principal
C) magnetic
D) A and B
E) B and C

Answer: D


27) Which quantum numbers must be the same for the orbitals that they designate to be degenerate in a many-electron system?

- A) n , l , m_l , and m_s
B) n and l only
C) n only
D) n , l , and m_l
E) m_s only


Answer: B

28) Which electron configuration represents a violation of the Pauli exclusion principle?


A)




B)




C)



D)




E)




Answer: A

29) Which electron configuration denotes an atom in its ground state?


A)




B)




C)



D)



E)



Answer: C

30) The lowest orbital energy is reached when the number of electrons with the same spin is maximized.

This statement describes _____.

- A) deBroglie hypothesis
- B) Hund's rule
- C) Planck's constant
- D) Heisenberg Uncertainty Principle
- E) Pauli Exclusion Principle

Answer: B

31) The wavelength of a photon that has an energy of 5.25×10^{-19} J is _____ m.

$$h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$$

- A) 2.64×10^6
- B) 3.79×10^{-7}
- C) 2.38×10^{23}
- D) 4.21×10^{-24}
- E) 3.79×10^7

Answer: B

32) The condensed electron configuration of silicon, element 14, is _____.

- A) $[\text{He}]2s^4$
- B) $[\text{He}]2s^4 2p^6$
- C) $[\text{He}]2s^6 2p^2$
- D) $[\text{Ne}]2p^{10}$
- E) $[\text{Ne}]3s^2 3p^2$

Answer: E

- 33) The $n = 5$ to $n = 3$ transition in the Bohr hydrogen atom corresponds to the _____ of a photon with a wavelength of _____ nm.

$$\frac{1}{\lambda} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \quad R_H = 1.096776 \times 10^7 \text{ m}^{-1}.$$

- A) absorption, 657
- B) emission, 657
- C) absorption, 1280
- D) emission, 389
- E) emission, 1280

Answer: E

- 34) In the following list, only _____ is not an example of matter.

- A) dust
- B) planets
- C) table salt
- D) elemental phosphorus
- E) light

Answer: E

- 35) A combination of sand, salt, and water is an example of a _____.

- A) pure substance
- B) solid
- C) heterogeneous mixture
- D) homogeneous mixture
- E) compound

Answer: C

- 36) Which one of the following has the element name and symbol correctly matched?

- A) Ag, silver
- B) Mg, manganese
- C) C, copper
- D) P, potassium
- E) Sn, silicon

Answer: A

- 37) Which of the following is not an exact number?

- A) The number of centimeters in an inch.
- B) The number of seconds in a ~~year~~. day
- C) The number of millimeters in a kilometer.
- D) The number of liters in a gallon.
- E) The number of grams in a kilogram.

Answer: D

- 38) The length of the side of a cube having a density of 12.6 g/ml and a mass of 7.65 g is _____ cm.

- A) 0.847
- B) 3.20
- C) 0.584
- D) 1.02
- E) 1.32

Answer: A

39) Osmium has a density of 22.6 g/cm^3 . The mass of a block of osmium that measures $1.01 \text{ cm} \times 0.233 \text{ cm} \times 0.648 \text{ cm}$ is _____ g.

- A) 3.45 B) 6.75×10^3 C) 6.75×10^{-3} D) 34.5 E) 148

Answer: A

40) The correct answer (reported to the proper number of significant figures) to the following is _____.

$$(2.05631)(6.9391136) / 12.59326 = \underline{\hspace{2cm}}$$

- A) 1.133064
B) 1.1361
C) 1.13306
D) 1.1330639
E) none of the above

Answer: C

41) The density of mercury is 13.6 g/cm^3 . The density of mercury is _____ kg/m^3 .

- A) 1.36×10^{-5} B) 1.36×10^{-4} C) 1.36×10^4 D) 1.36×10^{-2} E) 1.36×10^8

Answer: C

42) Consider the following selected postulates of Dalton's atomic theory:

- (i) Each element is composed of extremely small particles called atoms.
- (ii) Atoms are indivisible.
- (iii) Atoms of a given element are identical.
- (iv) Atoms of different elements are different and have different properties.

Which of the postulates is(are) no longer considered valid?

- A) (iii) only B) (iii) and (iv) C) (ii) only D) (ii) and (iii) E) (i) and (ii)


Answer: D

43) Which one of the following is not true concerning cathode rays?

- A) They travel in straight lines in the absence of electric or magnetic fields.
- B) They impart a negative charge to metals exposed to them.
- C) They originate from the negative electrode.
- D) They are made up of electrons.
- E) The characteristics of cathode rays depend on the material from which they are emitted.

Answer: E

44) Silver has two naturally occurring isotopes with the following isotopic masses:



The average atomic mass of silver is 107.8682 amu. The fractional abundance of the lighter of the two isotopes is _____.

- A) 0.75783 B) 0.24221 C) 0.48168 D) 0.51835 E) 0.90474

Answer: D

45) The average atomic weight of copper, which has two naturally occurring isotopes, is 63.5. One of the isotopes has an atomic weight of 62.9 amu and constitutes 69.1% of the copper isotopes. The other isotope has an abundance of 30.9%. The atomic weight (amu) of the second isotope is _____ amu.

- A) 28.1 B) 63.8 C) 64.1 D) 64.8 E) 63.2

Answer: D

46) Which formula/name pair is incorrect?

- A) $\text{Mg}(\text{MnO}_4)_2$ magnesium permanganate
B) $\text{Mn}(\text{NO}_3)_2$ manganese(II) nitrate
C) Mg_3N_2 magnesium nitrite
D) $\text{Mg}(\text{NO}_3)_2$ magnesium nitrate
E) $\text{Mn}(\text{NO}_2)_2$ manganese(II) nitrite

Answer: C

47) The correct name for K_2S is _____.

- A) potassium bisulfide
B) dipotassium sulfate
C) potassium disulfide
D) potassium sulfate
E) potassium sulfide

Answer: E

48) The correct name for HIO_2 is _____.

- A) iodic acid
B) periodic acid
C) hypoiodic acid
D) hydriodic acid
E) periodous acid

Answer: A

49) What is the molecular formula for 1-hexanol?

- A) $\text{C}_6\text{H}_{12}\text{OH}$ B) $\text{C}_7\text{H}_{13}\text{OH}$ C) $\text{C}_6\text{H}_{14}\text{OH}$ D) $\text{C}_7\text{H}_{14}\text{OH}$ E) $\text{C}_6\text{H}_{13}\text{OH}$

Answer: E

50) The name of the ionic compound RbBrO_4 is _____.

- A) rubidium bromate
B) rubidium hypobromate
C) rubidium bromide
D) rubidium perbromite
E) rubidium perbromate

Answer: E